



Data Article

Characteristics of soils in selected maize growing sites along altitudinal gradients in East African highlands



CrossMark

Elijah Njuguna ^{a,b}, Mary Gathara ^c, Stanley Nadir ^c, Sizah Mwalusepo ^{a,d}, David Williamson ^e, Pierre-Etienne Mathé ^f, Jackson Kimani ^a, Tobias Landmann ^a, Gerald Juma ^g, George Ong'amo ^g, Erastus Gatebe ^b, Bruno Le Ru ^{a,h}, Paul-andré Calatayud ^{a,h,*}

^a African Insect Science for Food and Health (icipe), P.O. Box 30772-00100, Nairobi, Kenya^b Jomo Kenyatta University of Agriculture and Technology (JKUAT), P.O. Box 62000, Nairobi, Kenya^c Kenya Forestry Research Institute (KEFRI), P.O. Box 20412-00200, Nairobi, Kenya^d Department of General Studies, Dar es Salaam Institute of Technology (DIT), P.O. Box 2958, Dar es Salaam, Tanzania^e Institut de Recherche pour le Développement (IRD), UMR-7153 LOCEAN, Université Pierre et Marie Curie, Centre IRD France Nord, 32 Avenue Henri-Varagnat, F-93143 Bondy cedex, France^f Aix-Marseille Université, CNRS, IRD, CEREGE UM34, Aix-en-Provence, France^g University of Nairobi, P.O. Box 30197-00100, Nairobi, Kenya^h UMR Laboratoire Evolution, Génomes, Comportement et Ecologie, groupe IRD, Diversité, Ecologie et Evolution des Insectes Tropicaux, UPR 9034, 22 CNRS, 91198 – Gif-sur-Yvette, France and Université de Paris-Sud, 91405-Orsay, France

ARTICLE INFO

Article history:

Received 16 June 2015

Received in revised form

24 August 2015

Accepted 25 August 2015

Available online 3 September 2015

Keywords:

Soil macro and micro elements

Organic carbon content

Soil texture

Soil pH

Cation exchange capacity

Electric conductivity

Water holding capacity

Leaching

ABSTRACT

Maize is the main staple crop in the East African Mountains. Understanding how the edaphic characteristics change along altitudinal gradients is important for maximizing maize production in East African Highlands, which are the key maize production areas in the region. This study evaluated and compared the levels of some macro and micro-elements (Al, Ca, Fe, K, Mg, Mn, Na and P) and other soil parameters (pH, organic carbon content, soil texture [i.e. % Sand, % Clay and % Silt], cation exchange capacity [CEC], electric conductivity [EC], and water holding capacity [HC]). Soil samples were taken from maize plots along three altitudinal gradients in East African highlands (namely Machakos Hills, Taita Hills and Mount Kilimanjaro) characterized by graded changes in climatic conditions. For all transects, pH, Ca, K and Mg decreased with the increase in altitude. In contrast, % Silt, organic carbon

* Corresponding author at: African Insect Science for Food and Health (icipe), P.O. Box 30772-00100, Nairobi, Kenya.
Tel.: +254 20 8632161; fax: +254 20 8632001/8632002.

E-mail address: pcalatayud@icipe.org (P.-a. Calatayud).